

INRNE-TH-97/08

September 1997

On the top-quark polarization and how to measure it

Ekaterina Christova

*Institute of Nuclear Research and Nuclear Energy
Boul. Tzarigradsko Chaussee 72, Sofia 1784, Bulgaria*

and

Dimitar Draganov

*Physics Department, Sofia University
Boul. James Baucher 5, Sofia 1126, Bulgaria* ABSTRACT

The top quark is expected to decay as a free particle with definite momentum and polarization. Here we consider the possibilities to obtain information about its polarization through the energy and angular distributions of the b-quarks from the decay of the top-quarks produced in e^+e^- annihilation. Analytic expressions and numerical estimates for different beam polarizations are derived.

1. The large mass of the top quark measured at Fermilab [1] allows to probe physics at high energies where new physics is expected to appear as well. Many papers treat this idea. That is why it becomes of basic interest to test first of all the Standard Model (S.M.) for processes with top quarks.

There is one basic difference of the top-quark as compared to the other quarks: due to its high mass, the top quark should decay before forming a hadronic bound state [2]. This means that the top quark will decay with a definite polarization, that can be measured through the distribution of its decay products, and that the theoretical predictions become more reliable free of the complications of hadronization. Measuring the top-quark polarization and comparing it to the theoretical predictions would present a clear test of the S.M.

All $t\bar{t}$ events contain a b and a $\bar{b}$ -quark. Different methods for detecting the $t\bar{t}$ events, based on tagging b -jets (for example via high $p_\perp$ leptons from $b \rightarrow c(i^\pm\nu)$ or via displaced vertices of b and $\bar{b}$ -decays) exist [3].

Here we discuss the possibilities to answer the questions whether the t -quark decays keeping its polarization and whether this polarization is the one predicted by S.M., measuring the energy and angular distributions of the b -quarks in the inclusive process:

$$e^+(q_l) + e^-(q_{\bar{l}}) \rightarrow t(p_t) + \bar{t}(p_{\bar{t}}) \rightarrow b(p_b) + X. \quad (1)$$

X stands for $W\bar{t}$. We assume that the initial beams are longitudinally polarized.

As the t -quark actually does not mix with other quarks, the considered decay mode $t \rightarrow bW$ is the only decay mode in S.M. This implies that the rate for production of b -quarks in (1) will be the same as the rate for production of t -quarks in e^+e^- annihilation.

Previously the t -quark polarization was considered in refs. [4], [5] and [6]. It was shown that the S.M. predicts a rather large polarization in the production plane of the t -quarks produced in the annihilation process of unpolarized e^+e^- . The possibilities to measure it through the angular distribution of the leptons from the decay mode $t \rightarrow bW \rightarrow bl\nu$ were considered in [4], [6] and [7].

Here we obtain analytic expressions for the energy and angular distributions of the b -quarks in the c.m.system of the sequential process (1). We follow the formalism of [8], in which the t -quark polarization enters explicitly. Different energy and angular asymmetries are defined. We show that measurements of the energy and angular distributions of the b -quarks would allow to determine the t -

quark polarization in the production plane. Numerical estimates for different beam polarizations and the dependence on the incertainties in m_t are presented.

2. Following the formalizm developed in [8], for the cross section $d\sigma_{\lambda\lambda'}$ of the sequential process (1), λ and λ' being the longitudinal polarizations of e^- and e^+ , we obtain:

$$\frac{d\sigma_{\lambda\lambda'}}{d\cos\theta d\Omega_b} = \left(\frac{d\sigma_{\lambda\lambda'}}{d\cos\theta d\Omega_b} \right)_0 \left(1 + \alpha_b m_t \frac{(\xi p_b)}{(p_t p_b)} \right). \quad (2)$$

All quantities in (2) are in the c.m.system, ξ is the polarization 4-vector of the t -quarks determined in the production process $e^+e^- \rightarrow t\bar{t}$, $\cos\theta$ is the production angle of the t -quarks (Z points the direction of e^-), Γ is the total decay width of the t -quark,

$$\alpha_b = \frac{m_t^2 - 2m_W^2}{m_t^2 + 2m_W^2} \quad (3)$$

and its value determines the sensitivity of the b-quarks to the polarization of the t -quarks, $(p_t p_b) = (m_t^2 - m_W^2)/2$,

$$\left(\frac{d\sigma_{\lambda\lambda'}}{d\cos\theta d\Omega_b} \right)_0 = \frac{\alpha_{em}^3}{32 \sin^2 \theta_w} |U_{tb}|^2 \frac{3\beta}{s} \frac{2m_W^2 + m_t^2}{m_W^2} \frac{E_b^2}{m_t \Gamma} N_{\lambda\lambda'} \quad (4)$$

is the S.M. cross section for totally depolarized t -quarks, $\sqrt{s}$ is the c. m. energy, $\beta = \sqrt{1 - 4m_t^2/s}$, U_{tb} is the corresponding element of the Kobayashi-Maskawa matrix, $N_{\lambda\lambda'}$ equals

$$N_{\lambda\lambda'} = \left(1 + \beta^2 \cos^2 \theta \right) F_1 + \frac{4m_t^2}{s} F_2 + 2\beta \cos \theta F_3, \quad (5)$$

where

$$F_i = (1 - \lambda\lambda') F_i^0 + (\lambda - \lambda') G_i^0. \quad (6)$$

Here

$$\begin{aligned} F_{1,2}^0 &= \left(\frac{2}{3} \right)^2 + h_Z^2 (c_V^2 + c_A^2) (g_V^2 \pm g_A^2) - \frac{4}{3} h_Z c_V g_V \\ G_{1,2}^0 &= 2h_Z^2 c_V c_A (g_V^2 \pm g_A^2) - \frac{4}{3} h_Z c_A g_V, \\ F_3^0 &= 4h_Z c_A g_A \left(h_Z c_V g_V - \frac{1}{3} \right), G_3^0 = 2h_Z^2 (c_V^2 + c_A^2) g_V g_A - \frac{4}{3} h_Z c_V g_A, \end{aligned} \quad (7)$$

E_b is the energy of the b-quarks in the c.m.system: $E_b = (p_t p_b)/[E(1 - \beta \cos \theta_{tb})]$, E is the energy of the initial leptons, $\cos \theta_{tb}$ is the decay angle between the directions of the t and b-quarks, $c_V = -1/2 + 2 \sin^2 \theta_W$, $c_A = 1/2$ and $g_V = 1/2 - 4/3 \sin^2 \theta_W$, $g_A = -1/2$ are the S.M. couplings of Z^0 to the electron and the top-quark respectively, $h_Z = s/[\sin^2 2\theta_w(s - m_Z^2)]$.

3. Now we obtain the expression for the polarization 4-vector ξ^α of the top-quarks that are produced in e^+e^- annihilation with longitudinally polarized initial leptons in tree-level approximation of the S.M.

As by definition $(\xi p_t) = 0$, ξ_α in general (in tree level) can be decomposed along two independent 4-vectors, orthogonal to p_t , that lay in the production plane. We choose them to be Q_l and $Q_{\bar{l}}$:

$$Q_l^\alpha = q_l^\alpha - \frac{p_t q_l}{m_t^2} \cdot p_t^\alpha, \quad Q_{\bar{l}}^\alpha = q_{\bar{l}}^\alpha - \frac{p_t q_{\bar{l}}}{m_t^2} \cdot p_t^\alpha. \quad (8)$$

Then we write:

$$(\xi)^\alpha = P_l(Q_l)^\alpha + P_{\bar{l}}(Q_{\bar{l}})^\alpha. \quad (9)$$

Using the method of [8] we obtain:

$$P_{l,\bar{l}}(\theta) = \pm \frac{2m_t}{s} \{ (1 \mp \beta \cos \theta)(G_1 \mp G_3) + (1 \pm \beta \cos \theta)G_2 \} / N_{\lambda\lambda'} \quad (10)$$

Here

$$G_i = (1 - \lambda\lambda')G_i^0 + (\lambda - \lambda')F_i^0, \quad (11)$$

F_i^0 and G_i^0 are given by (7).

Having the explicit expression for ξ^α , from (2) one readily obtains the energy and angular distributions of the b-quarks.

There are two independent components of the polarization vector of the top in tree level of the S.M. Thus we need at least two independent measurements in order to get information about ξ^α . As it is evident from (2) the polarization of the t-quarks 1) introduces a new dependence on $\cos \theta_b$, not only through $\cos \theta_{tb}$, and 2) changes the behaviour on $\cos \theta_{tb}$, or equivalently on E_b . We shall explore both to get information about ξ .

4. For the energy spectrum of the b -quarks in the c.m.s. we obtain:

$$\frac{d\sigma}{dE_b} = \left(\frac{d\sigma}{dE_b} \right)_0 \left[1 + \frac{4\alpha_b G_3}{(3 + \beta^2)F_1 + 3(1 - \beta^2)F_2} \left(1 - \frac{4m_t^2}{m_t^2 - m_W^2} \frac{E_b}{\sqrt{s}} \right) \right] \quad (12)$$

where

$$\begin{aligned} \left(\frac{d\sigma}{dE_b} \right)_0 &= \frac{\alpha_{em}^3}{8 \sin^2 \theta_w s \sqrt{s}} \pi |U_{tb}|^2 \frac{m_t^2 + 2m_W^2}{m_W^2} \frac{m_t^2 - m_W^2}{m_t \Gamma} \times \\ &\times [(3 + \beta^2)F_1 + 3(1 - \beta^2)F_2] \end{aligned} \quad (13)$$

is the energy spectrum for unpolarized t -quarks. Thus, if the t -quarks decay depolarized, the cross section for a given energy E_b would not depend on the energy of the b -quark. The polarization of the decaying t -quarks turns this constant behaviour into a linear E_b -behaviour, the decline determined by G_3 . This allows to form different asymmetries, whose nonzero value would imply a nonzero polarization of the t -quarks.

If $N_{\lambda\lambda'}(E_b)$ is the number of b -quarks with energy E_b , then the ratio $R(\Delta E_b)$:

$$R_{\lambda\lambda'}(\Delta E_b) = \frac{N_{\lambda\lambda'}(E_2) - N_{\lambda\lambda'}(E_1)}{N_{\lambda\lambda'}(E_2) + N_{\lambda\lambda'}(E_1)}, \quad \Delta E_b = E_2 - E_1 \quad (14)$$

would be proportional to the top-polarization. As the energy spectrum exhibits a linear dependence on the polarization, $R(\Delta E_b)$ would reach its maximum if measurements are fulfilled at the ends of the spectrum: $E_2 = E_{max}$, $E_1 = E_{min}$. Then we obtain:

$$R_{\lambda\lambda'}(E_{max} - E_{min}) = \frac{-4\alpha_b \beta G_3}{(3 + \beta^2)F_1 + 3(1 - \beta^2)F_2}. \quad (15)$$

One can form different integral asymmetries comparing the number of b -quarks in two energy intervals with the same length. We consider the following asymmetry, whose nonzero value would be an indication of polarized decaying t -quarks.:

$$\mathcal{R}_{\lambda\lambda'} = \frac{N_{\lambda\lambda'}(E_b > \mathcal{E}_0) - N_{\lambda\lambda'}(E_b < \mathcal{E}_0)}{N_{\lambda\lambda'}(E_b > \mathcal{E}_0) + N_{\lambda\lambda'}(E_b < \mathcal{E}_0)} = \frac{-2\alpha_b \beta G_3}{(3 + \beta^2)F_1 + 3(1 - \beta^2)F_2}. \quad (16)$$

where $\mathcal{E}_0$ is the mean energy: $\mathcal{E}_0 = (E_{min} + E_{max})/2 = \sqrt{s}(m_t^2 - m_W^2)/(4m_t^2)$, $N_{\lambda\lambda'}(E_b > \mathcal{E}_0)$ and $N_{\lambda\lambda'}(E_b < \mathcal{E}_0)$ are the number of b -quarks with energy $E_b > \mathcal{E}_0$ and $E_b < \mathcal{E}_0$ respectively.

5. Integrating (2) over θ and φ_b we obtain the $\cos \theta_b$ -distribution of the b -quarks in the c.m.system. We write it in a form that keeps in evidence the dependence on the beam polarization too:

$$\begin{aligned} \frac{d\sigma_{\lambda\lambda'}}{d(\cos \theta_b)} &= \frac{3\alpha_{em}^3}{64 s \sin^2 \theta_w} \pi |U_{tb}|^2 \frac{(m_t^2 - m_W^2)^2 (m_t^2 + 2m_W^2)}{m_t^3 m_W^2 \Gamma} \times \\ &\times \left\{ (1 - \lambda\lambda') [C_0 + C_1 \cos \theta_b + C_2 \cos^2 \theta_b] \right. \\ &\quad \left. + (\lambda - \lambda') [O_0 + O_1 \cos \theta_b + O_2 \cos^2 \theta_b] \right\} \end{aligned} \quad (17)$$

where

$$\begin{aligned} C_0 &= F_1^0 \left(2\beta^3 + (1 - \beta^2) \ln \frac{1 + \beta}{1 - \beta} \right) + 2\beta(1 - \beta^2) F_2^0 \\ &\quad - \alpha_b(1 - \beta^2) G_3^0 \left(2\beta - \ln \frac{1 + \beta}{1 - \beta} \right) \end{aligned} \quad (18)$$

$$\begin{aligned} C_1 &= 2F_3^0 \left(2\beta - (1 - \beta^2) \ln \frac{1 + \beta}{1 - \beta} \right) \\ &\quad + 2\alpha_b(1 - \beta^2) \left[\beta(G_1^0 - G_2^0) - G_1^0 \ln \frac{1 + \beta}{1 - \beta} \right] \end{aligned} \quad (19)$$

$$\begin{aligned} C_2 &= F_1^0 \left(2\beta(3 - 2\beta^2) - 3(1 - \beta^2) \ln \frac{1 + \beta}{1 - \beta} \right) \\ &\quad + 3\alpha_b(1 - \beta^2) G_3^0 \left(2\beta - \ln \frac{1 + \beta}{1 - \beta} \right) \end{aligned} \quad (20)$$

$$O_i = C_i (F_i^0 \longleftrightarrow G_i^0) \quad (21)$$

As it is evident from (18) - (20), the polarization of the t -quarks does not introduce any new type of $\cos \theta_b$ -dependence. Thus, the angular-polarization asymmetries, contrary to the considered energy asymmetries, would be nonzero for both polarized and depolarized t -quarks, but differ numerically.

Let $\sigma_{\lambda\lambda'}^F$ and $\sigma_{\lambda\lambda'}^B$ denote the number of b -quarks produced in the forward and backward hemispheres respectively:

$$\sigma_{\lambda\lambda'}^F = 2\pi \int_{\theta_0}^{\pi/2} \left(\frac{d\sigma_{\lambda\lambda'}}{d \cos \theta_b} \right) \sin \theta_b d\theta_b, \quad \sigma_{\lambda\lambda'}^B = 2\pi \int_{\pi/2}^{\pi - \theta_0} \left(\frac{d\sigma_{\lambda\lambda'}}{d \cos \theta_b} \right) \sin \theta_b d\theta_b$$

θ_0 is determined by the experimental set up. The forward-backward asymmetries that we shall consider are sensitive to the combinations $\left[\beta(G_1^0 - G_2^0) - G_1^0 \ln \frac{(1+\beta)}{(1-\beta)} \right]$

and $\left[\beta(F_1^0 - F_2^0) - F_1^0 \ln \frac{(1+\beta)}{(1-\beta)}\right]$ that enter C_1 and O_1 . The other contribution to ξ^α , determined by G_3^0, F_3^0 that enter $C_{0,2}, O_{0,2}$ can be measured by the energy spectrum, as shown above. Thus, measuring the energy and angular distributions of the b -quarks, with either polarized or unpolarized initial beams we obtain information about two independent contributions to the polarization of the t -quark.

We consider the following angular-polarization b -jet asymmetries:

i) The forward-backward asymmetry $A_{\lambda\lambda'}^{FB}$, that measures the difference between the b -quarks in the forward and backward hemispheres:

$$A_{\lambda\lambda'}^{FB} = \frac{\sigma_{\lambda\lambda'}^F - \sigma_{\lambda\lambda'}^B}{\sigma_{\lambda\lambda'}^F + \sigma_{\lambda\lambda'}^B} \quad (22)$$

From (17) for $A_{\lambda\lambda'}^{FB}$ we obtain:

$$A_{\lambda\lambda'}^{FB} = \frac{(1 - \lambda\lambda') C_1 + (\lambda - \lambda') O_1}{(1 - \lambda\lambda') (2 C_0 + 2 C_2/3) + (\lambda - \lambda') (2 O_0 + 2 O_2/3)} \quad (23)$$

Our expressions (17) and (23) for the $\cos \theta_b$ -distribution and for the forward-backward asymmetry (23) for unpolarized initial leptons ($\lambda = \lambda' = 0$) coincide with those obtained in [6], where they were first derived using the technique of ref [9].

ii) The polarization angular asymmetry $\mathcal{A}^{FB}$, that compares the b -quarks in the forward and backward hemispheres with opposite beam polarizations:

$$\mathcal{A}^{FB} = \frac{\sigma_{\lambda\lambda'}^F - \sigma_{-\lambda-\lambda'}^B}{\sigma_{\lambda\lambda'}^F + \sigma_{-\lambda-\lambda'}^B} = \frac{(1 - \lambda\lambda') C_1 + (\lambda - \lambda') (2 O_0 + 2 O_2/3)}{(1 - \lambda\lambda') (2 C_0 + 2 C_2/3) + (\lambda - \lambda') O_1} \quad (24)$$

iii) The polarization asymmetry $\mathcal{B}_{LR}^{FB}$ that singles out O_1 and C_1 only:

$$\mathcal{B}^{FB} = \frac{1 - \lambda\lambda'}{\lambda - \lambda'} \frac{(\sigma_{\lambda\lambda'}^F - \sigma_{\lambda\lambda'}^B) - (\sigma_{-\lambda-\lambda'}^F - \sigma_{-\lambda-\lambda'}^B)}{(\sigma_{\lambda\lambda'}^F - \sigma_{\lambda\lambda'}^B) + (\sigma_{-\lambda-\lambda'}^F - \sigma_{-\lambda-\lambda'}^B)} = \frac{O_1}{C_1} \quad (25)$$

Note that $\mathcal{B}^{FB}$ is defined for polarized beams only and is independent on the beam polarization, $\mathcal{B}^{FB}(\lambda' = \lambda = 0) = 0$.

The above asymmetries contain also the combinations $2 C_0 + 2 C_2/3$ and $2 O_0 + 2 O_2/3$. These combinations enter the total cross section and are independent on the t -quark polarization.

6. We have estimated the defined asymmetries at $\sqrt{s} = 500$ GeV assuming $\lambda' = -\lambda$. We examine their sensitivity to the uncertainties in m_t for different beam polarizations.

Observation of a linear E_b -dependence of the b -quark spectrum in stead of a constant one, or equivalently non-zero values of R or $\mathcal{R}$ would be possible indications for polarized decaying t -quarks. Their values are determined by G_3 . On Fig. 1 we show the dependence of $R(E_{max} - E_{min})$ on the beam polarization λ . R depends strongly on λ - it varies from $\approx 6\%$ for $\lambda = 0$ up to $\approx 17\%$ for $\lambda = \pm 0.5$. As our analysis showed it is actually independent on m_t . The integral asymmetry $\mathcal{R}$ should be two times less, as evident from (16).

The sensitivity of the angular asymmetries A^{FB} , $\mathcal{A}^{FB}$ and $\mathcal{B}^{FB}$ to ξ is determined by the difference between their values A^{FB} , $\mathcal{A}^{FB}$ and $\mathcal{B}^{FB}$ with polarized decaying t -quarks - eqs. (23), (24) and (25), and the corresponding values $\bar{A}^{FB}$, $\bar{\mathcal{A}}^{FB}$ and $\bar{\mathcal{B}}^{FB}$ with depolarized decaying t -quarks ($\alpha_b = 0$). It is determined by G_1 and G_2 .

On Fig. 2, the values of A^{FB} and $\bar{A}^{FB}$ are shown as functions of λ . The difference $A^{FB} - \bar{A}^{FB}$ is strongly dependent on λ . For unpolarized beams one needs around 5% accuracy of measurements, while for $\lambda = \pm 0.5$, an accuracy of 10% would be enough to disentangle A^{FB} from $\bar{A}^{FB}$. Note that the value of this asymmetry is rather large – even for unpolarized t -quarks it is about 20%.

On Table 1 we draw attention that for unpolarized beams the uncertainties in m_t might not allow to make clear predictions about the t -polarization – the m_t -dependence can mimic the ξ -dependence of A^{FB} . For example, at $\lambda = 0$ we have $\bar{A}^{FB}(m_t = 165\text{GeV}) = A^{FB}(m_t = 185\text{GeV})$. The beam polarization would allow to distinguish the decay of polarized and depolarized t -quarks independently on the uncertainties in m_t : at $\lambda = -0.5$, $A^{FB} - \bar{A}^{FB} \geq 10\%$ for any value of m_t .

Fig.3 shows the dependence of $\mathcal{A}^{FB}$ and $\bar{\mathcal{A}}^{FB}$ on λ for $m_t = 175\text{GeV}$. It is seen that the difference $\mathcal{A}^{FB} - \bar{\mathcal{A}}^{FB}$ does not exceed 5%. As shown in Table 1 the uncertainties in m_t can smear this difference and higher precision for m_t would be necessary to use this asymmetry for to determination of the t -polarization.

The polarization asymmetry $\mathcal{B}^{FB}$ appears to be the most sensitive one to t -quark polarization - the difference $\mathcal{B}^{FB} - \bar{\mathcal{B}}^{FB}$ is of about 40% and it is not reduced by the uncertainties in m_t (see Table 1). The magnitude of the asymmetry is also quite large: for $m_t = 175\text{ GeV}$, $\mathcal{B}^{FB} \approx -0.74$.

7. We are thankful to Serguey Petcov for the helpful remarks and discussion. This work was supported by the Bulgarian National Science Foundation, Grant 510

m_t	$A^{FB}(0)$	$\bar{A}^{FB}(0)$	$A^{FB}(0.5)$	$\bar{A}^{FB}(0.5)$	$A^{FB}(-0.5)$	$\bar{A}^{FB}(-0.5)$
165	0.29	0.25	0.22	0.29	0.35	0.23
175	0.28	0.22	0.16	0.24	0.34	0.21
185	0.25	0.19	0.12	0.20	0.34	0.18
m_t	$\mathcal{A}_{LR}^{FB}(0.5)$	$\bar{\mathcal{A}}_{LR}^{FB}(0.5)$	$\mathcal{A}_{LR}^{FB}(-0.5)$	$\bar{\mathcal{A}}_{LR}^{FB}(-0.5)$	$\mathcal{B}^{FB}$	$\bar{\mathcal{B}}^{FB}$
165	0.00	-0.05	0.52	0.52	-0.63	-0.28
175	-0.03	-0.08	0.50	0.50	-0.74	-0.28
185	-0.06	-0.12	0.47	0.47	-0.86	-0.28

Table 1: The dependence of the angular asymmetries on m_t , in brackets is the beam polarization $\lambda = 0; \pm 0.5$, $\mathcal{A}^{FB}(0) = A^{FB}(0)$, $\mathcal{B}^{FB}(0) = 0$

References

- [1] CDF Collaboration, F.Abe et al, *Phys.Rev.Lett.* **74** (1995) 2626
D0 Collaboration, S. Abachi et al, *Phys.Rev.Lett.* **74** (1995) 2632
- [2] I. Bigi, V. Dokshitzer, V. Khoze, J. Kuhn and P.M. Zerwas, *Phys. Lett* **B181** (1986) 157
- [3] G. Bagliesi et al. e^+e^- Collisions at 500 GeV: The Physical Potential, Part A, DESY 92-123A (1992),p.327
- [4] R.H. Dalitz and G.R. Goldstein, *Phys. Rev.***D45** (1992) 1531
- [5] G.Kane, Landinski and C.P. Yuan, *Phys. Rev.* **D45** (1992) 124
- [6] T. Arens and L.M. Sehgal, *Nucl. Phys.***B393** (1993) 46 and references therein
- [7] A. Czarnecki, M.Jazabeck and J.H. Kuhn, *Nucl. Phys.* **B351** (1991) 70

[8] S.M.Bilenky, E.Christova, N.Nedelcheva, *Bulg.J.Phys.***13** (1986) 283

[9] S. Kawasaki, T. Shirafuji and S Tsai, *Progress of Theor. Phys.* **49** (1973) 1656

Figure Captions

Fig. 1: The energy asymmetry $R(E_{max} - E_{min})$ as a function of the beam polarization λ , $m_t = 175$ GeV.

Fig. 2: The asymmetry A^{FB} for polarized (full curve) and depolarized (dashed curve) t -quarks as a function of λ , $m_t=175$ GeV.

Fig. 3: The asymmetry $\mathcal{A}^{FB}$ for polarized (full curve) and depolarized (dashed curve) t -quarks as a function of λ , $m_t=175$ GeV.




Fig. 1




Fig. 2




Fig. 3